


Radius of an atom
 $1 \times 10^{-10} \text{m}$



Electrons gained
Negative ion

Electrons lost
Positive ion



Atom	Same number of protons and electrons
Ion	Unequal number of electrons to protons
Mass number	Number of protons <u>and</u> neutrons
Atomic number	Number of protons

Particle	Charge	Size	Found
Neutron	None	1	In the nucleus
Proton	+	1	
Electron	-	Tiny	

Isotope	${}^6_3\text{Li}$		${}^7_3\text{Li}$	
Different forms of an element with the same number of protons but different number of neutrons				

Discovery of the nucleus

Democritus	Suggested idea of atoms as small spheres that cannot be cut.
J J Thomson (1897)	Discovered electrons— emitted from surface of hot metal. Showed electrons are negatively charged and that they are much less massive than atoms.
Thomson (1904)	Proposed 'plum pudding' model – atoms are a ball of positive charge with negative electrons embedded in it.
Geiger and Marsden (1909)	Directed beam of alpha particles (He^{2+}) at a thin sheet of gold foil. Found some travelled through, some were deflected, some bounced back.
Rutherford (1911)	Used above evidence to suggest alpha particles deflected due to electrostatic interaction between the very small charged nucleus, nucleus was massive. Proposed mass and positive charge contained in nucleus while electrons found outside the nucleus which cancel the positive charge exactly.
Bohr (1913)	Suggested modern model of atom – electrons in circular orbits around nucleus, electrons can change orbits by emitting or absorbing electromagnetic radiation. His research led to the idea of some particles within the nucleus having positive charge; these were named protons.
Chadwick (1932)	Discovered neutrons in nucleus – enabling other scientists to account for mass of atom.

Atom structure

Atoms and Isotopes

Atoms and Nuclear Radiation

Radioactive decay	Unstable atoms randomly emit radiation to become stable
Detecting	Use Geiger Muller tube
Unit	Becquerel
Ionisation	All radiation ionises

Decay	Emitted from nucleus	Changes in mass number and atomic number	
Alpha (α)	Helium nuclei (${}^4_2\text{He}$)	-4	-2
Beta (β)	Electron (${}^0_{-1}\text{e}$)	0	+1
Gamma (γ)	Electromagnetic wave	0	0
Neutron	Neutron	-1	0

Contamination	Unwanted presence of radioactive atoms
Irradiation	Person is in exposed to radioactive source

AQA ATOMIC STRUCTURE

PHYSICS ONLY: Hazards and uses of Radioactive emissions and of background radiation

Half life
The time taken to lose half of its initial radioactivity

Sievert	Unit measuring dose of radiation
Background	Constant low level environmental radiation, e.g. from nuclear testing, nuclear power, waste


Uses	Different isotopes have different half lives	Short half-lives used in high doses, long half lives used in low doses.
Tracers	Used within body	Isotope with short half life injected, allowed to circulate and collect in damaged areas. PET scanner used to detect emitting radiation. Must be beta or gamma as alpha does not penetrate the body.
Radiation therapy	Used to treat illnesses e.g. cancer	Cancer cells killed by gamma rays. High dose used to kill cells. Damage to healthy cells prevented by focussed gamma ray gun.

Fuel rods	Made of U-238, 'enriched' with U-235 (3%). Long and thin to allow neutrons to escape, hitting nuclei.
Control rods	Made of Boron. Controls the rate of reaction. Boron absorbs excess neutrons.
Concrete	Neutrons hazardous to humans – thick concrete shield protects workers.

Nuclear fission and fusion

PHYSICS ONLY: Nuclear energy

Nuclear fission	One large unstable nucleus splits to make two smaller nuclei	Neutron hits U-235 nucleus, nucleus absorbs neutron, splits emitting two or three neutrons and two smaller nuclei. Process also releases energy.	Process repeats, chain reaction formed
Nuclear fusion	Two small nuclei join to make one larger nucleus	Difficult to do on Earth – huge amounts of pressure and temperature needed.	Occurs in stars



Decay	Range in air	Ionising power	Penetration power
Alpha	Few cm	Very strong	Stopped by paper
Beta	Few m	Medium	Stopped by Aluminium
Gamma	Great distances	Weak	Stopped by thick lead

